## Update website feedfoodtransfer.nl

## Last update March 21, 2022.

Below, a summary of features added and issues addressed on <u>feedfoodtransfer.nl</u>, since its release on January 27 2020, is provided. Changes are given per transfer model, starting with the most recent changes.

Feature	Description	
General website		
Update proclaimer (27-05-2020)	Addition of following sentence "To evaluate compliance with legal standards, the results obtained with the simulation of the transfer of the contaminant from feed to animal products should be confirmed by analytical results".	
Transfer model aflatoxin B1	L - dairy cow	
Version 1.1 (02-03-2021)	<ul> <li>Changes were made to improve the user-friendliness of the model including:</li> <li>The daily milk production can now be changed</li> <li>A warning will be given when negative values are entered in the input fields</li> <li>Results for milk (aflatoxin M1) and results for muscle, kidney and milk (aflatoxin B1 and M1) can be displayed simultaneously.</li> <li>Graph functions of Plotly are added.</li> <li>The report now contains the proclaimer, the model version used, the date, all input data and the output as visible online when the report is made.</li> </ul>	
	The manual is updated to comply with version 1.1.	
Version 1.0 (27-01-2020)	Release	
Transfer model dioxins (tot	al TEQ) - dairy cow	
Version 1.1 (02-03-2021)	<ul> <li>Body fat concentrations were incorrectly calculated in version 1.0. This is corrected in version 1.1.</li> <li>Changes were made to improve the user-friendliness of the model including:         <ul> <li>A warning will be given when negative values are entered in the input fields.</li> <li>Results for milk fat and body fat can be displayed simultaneously.</li> <li>Graph functions of Plotly are added.</li> <li>The report now contains the proclaimer, the model version used, the date, all input data and the output as visible online when the report is made.</li> </ul> </li> <li>The manual is updated to comply with version 1.1.</li> </ul>	
Version 1.0	Release	
(27-01-2020)		
Transfer model PFOS / PFOA – dairy cow		
Version 2.0 (21-03-2022)	The transfer model is updated and can now also be used to calculate the transfer of PFOA. The manual is updated to comply with version 2.0.	

	T
Version 1.1	A warning will be given when negative values are entered in
(02-03-2021)	the input fields
	Results for a lumped compartment based on the weighed
	sum of liver, kidney and muscle weight are now visible and
	can be displayed simultaneously with the results for blood
	and milk.
.,	The manual is updated to comply with version 1.1.
Version 1.0	Release
(01-12-2021)	
Transfer model cadmium -	
Version 1.0	Release
(27-01-2020)	
Transfer model dioxins (to	
Version 2.0	The transfer model is updated and a default time-dependent intake
(21-03-2022)	(in kg 88% dry matter per day) starting at day 28 (compound feed) or
	at day 66 (wet mixes) is available. The user can still use a specific
	feeding regime instead of this default regime.
	In addition changes were made to improve the user-friendliness of
	the model including:
	A warning will be given when negative values are entered in
	the input fields.
	Results for body fat and central compartment and the feed
	intake curve can be displayed simultaneously.
	<ul> <li>Graph functions of Plotly are added.</li> </ul>
	<ul> <li>The report now contains the proclaimer, the model version</li> </ul>
	used, the date, all input data and the output as visible online
	when the report is made.
	The manual is updated to comply with version 2.0.
Version 1.0	Release
(27-01-2020)	
Transfer model dioxins (to	tal TEQ) – laying hen
Version 1.1	For parameters y, F <sub>abs</sub> and k a set of optimal solutions can only be
(21-03-2022)	defined based on the upper and lower boundaries of allowable
	parameter values, Parameters y, F <sub>abs</sub> and k were set to the boundary
	solution Fabs,min; $k_{min} = 0$ ; $y_{max}$ ; $W_{fmin}$ ). This has basically no
	consequences for the estimations of the transfer model.
Version 1.0	Release
(01-12-2020)	
	ngener-specific) – laying hen
Version 1.3	For parameters y, F <sub>abs</sub> and k a set of optimal solutions can only be
(21-03-2022)	defined based on the upper and lower boundaries of allowable
	parameter values, Parameters y, F <sub>abs</sub> and k were set to the boundary
	solution Fabs,min; $k_{min} = 0$ ; $y_{max}$ ). This has basically no consequences
	for the estimations of the transfer model.
Version 1.2	The unit for the contaminant concentrations in feed in the
(02-03-2021)	downloadable xlsx.file is corrected
Version 1.1	Changes were made to improve the user-friendliness of the model
(01-12-2020)	including:
	<ul> <li>The CVS template for entering the contaminant</li> </ul>
	concentrations in feed is replaced by an xls.file

	<ul> <li>A warning will be given when negative values are entered in the input fields</li> <li>The option to insert a regulatory limit for body fat in addition to egg fat is added.</li> <li>The input fields for the congeners are now listed in separate tabs for PCD/F and dl-PCB's.</li> <li>Results for egg yolk fat and body fat can be displayed simultaneously.</li> <li>Results for all congeners can be displayed simultaneously.</li> <li>Graph functions of Plotly are added.</li> <li>The report now contains the proclaimer, the model version used, the date, all input data and the output as visible online when the report is made.</li> <li>The manual is updated to comply with version 1.1.</li> </ul>
Version 1.0 (27-01-2020)	Release